


Journal Pre-proof

Predictors of Patient Satisfaction in Spine Surgery: A Systematic Review

Brandon M. Lehrich, BS, Khodayar Goshtasbi, BS, Nolan J. Brown, BS, Shane Shahrestani, MS, Brian V. Lien, MS, Seth C. Ransom, BS, Ali R. Tafreshi, MD, Ryan C. Ransom, MD, PhD, Alvin Y. Chan, MD, Luis D. Diaz-Aguilar, MD, Ronald Sahyouni, MD, MS, PhD, Martin H. Pham, MD, Joseph A. Osorio, MD, PhD, Michael Y. Oh, MD



PII: S1878-8750(20)32510-9

DOI: <https://doi.org/10.1016/j.wneu.2020.11.125>

Reference: WNEU 16396

To appear in: *World Neurosurgery*

Received Date: 28 August 2020

Revised Date: 20 November 2020

Accepted Date: 21 November 2020

Please cite this article as: Lehrich BM, Goshtasbi K, Brown NJ, Shahrestani S, Lien BV, Ransom SC, Tafreshi AR, Ransom RC, Chan AY, Diaz-Aguilar LD, Sahyouni R, Pham MH, Osorio JA, Oh MY, Predictors of Patient Satisfaction in Spine Surgery: A Systematic Review, *World Neurosurgery* (2020), doi: <https://doi.org/10.1016/j.wneu.2020.11.125>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc.

Predictors of Patient Satisfaction in Spine Surgery: A Systematic Review

Brandon M. Lehrich, BS^{1*}; Khodayar Goshtasbi, BS^{2*}; Nolan J. Brown, BS^{3*}; Shane Shahrestani, MS^{4,5}; Brian V. Lien, MS³; Seth C. Ransom, BS⁶; Ali R. Tafreshi, MD⁷; Ryan C. Ransom, MD, PhD⁸; Alvin Y. Chan, MD³; Luis D. Diaz-Aguilar, MD⁹; Ronald Sahyouni, MD, MS, PhD⁹; Martin H. Pham, MD⁹; Joseph A. Osorio, MD, PhD⁹; Michael Y. Oh, MD³

¹Medical Scientist Training Program, University of Pittsburgh and Carnegie Mellon University, Pittsburgh, PA, USA

²School of Medicine, University of California, Irvine, Irvine, CA, USA

³Department of Neurosurgery, University of California, Irvine, Irvine, CA, USA

⁴Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁵Department of Medical Engineering, California Institute of Technology, Pasadena, CA, USA

⁶School of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA

⁷Department of Neurological Surgery, Geisinger Health System, Danville, PA, USA

⁸Department of Neurological Surgery, Mayo Clinic, Rochester, MN, USA

⁹Department of Neurological Surgery, University of California, San Diego, La Jolla, CA, USA

*These authors contributed equally to this manuscript.

Short running title: Spine Patient Satisfaction

Source of funding: Research reported in this publication was in part supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number T32GM008208 to BML. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Financial interests: None.

Disclosure: The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

Level of Evidence: NA

****Corresponding author:**

Brandon M. Lehrich, BS

Medical Scientist Training Program

University of Pittsburgh and Carnegie Mellon University

M246 Scaife Hall

3550 Terrace Street

Pittsburgh, PA 15232

Phone: (412) 648-2324

Email: brandon.lehrich@pitt.edu

Predictors of Patient Satisfaction in Spine Surgery: A Systematic Review

Brandon M. Lehrich, BS^{1*}; Khodayar Goshtasbi, BS^{2*}; Nolan J. Brown, BS^{3*}; Shane Shahrestani, MS^{4,5}; Brian V. Lien, MS³; Seth C. Ransom, BS⁶; Ali R. Tafreshi, MD⁷; Ryan C. Ransom, MD, PhD⁸; Alvin Y. Chan, MD³; Luis D. Diaz-Aguilar, MD⁹; Ronald Sahyouni, MD, MS, PhD⁹; Martin H. Pham, MD⁹; Joseph A. Osorio, MD, PhD⁹; Michael Y. Oh, MD³

¹Medical Scientist Training Program, University of Pittsburgh and Carnegie Mellon University, Pittsburgh, PA, USA

²School of Medicine, University of California, Irvine, Irvine, CA, USA

³Department of Neurosurgery, University of California, Irvine, Irvine, CA, USA

⁴Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

⁵Department of Medical Engineering, California Institute of Technology, Pasadena, CA, USA

⁶School of Medicine, University of Arkansas for Medical Sciences, Little Rock, AR, USA

⁷Department of Neurological Surgery, Geisinger Health System, Danville, PA, USA

⁸Department of Neurological Surgery, Mayo Clinic, Rochester, MN, USA

⁹Department of Neurological Surgery, University of California, San Diego, La Jolla, CA, USA

*These authors contributed equally to this manuscript.

Short running title: Spine Patient Satisfaction

Source of funding: Research reported in this publication was in part supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number T32GM008208 to BML. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Financial interests: None.

Disclosure: The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper.

Level of Evidence: NA

****Corresponding author:**

Brandon M. Lehrich, BS

Medical Scientist Training Program

University of Pittsburgh and Carnegie Mellon University

M246 Scaife Hall

3550 Terrace Street

Pittsburgh, PA 15232

Phone: (412) 648-2324

Email: brandon.lehrich@pitt.edu

Abstract

Background: Recently, there has been increased interest in patient satisfaction measures such as Press Ganey and Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) surveys. This systematic review aims to analyze the available spine surgery literature to evaluate factors predictive of patient satisfaction as measured by these surveys.

Methods: A thorough literature search was performed in PubMed/MEDLINE, Google Scholar, and Cochrane databases. All English-language articles from database inception to July 2020 were screened for study inclusion according to PRISMA guidelines.

Results: Twenty-four of the 1,899 published studies were included for qualitative analysis. There has been a statistically significant increase in the number of publications across years ($p=0.04$). Overall, the studies evaluated the relationship between patient satisfaction and patient demographics (71%), pre- and intra-operative clinical factors (21%), and post-operative factors (33%). Top positive predictors of patient satisfaction were patient and nursing/medical staff relationship ($n=4$; 17%), physician-patient relationship ($n=4$; 17%), managerial oversight of received care ($n=3$; 13%), same sex/ethnicity between patient and physician ($n=2$; 8%), and older age ($n=2$; 8%). Top negative predictors of patient satisfaction were high Charlson Comorbidity index/high disability/worse overall health functioning ($n=7$; 29%), increased length of hospital stay ($n=4$; 17%), high rating for pain/complications/readmissions ($n=4$; 17%), and psychosocial factors ($n=3$; 13%).

Conclusions: There is heterogeneity in terms of different factors, both clinical and non-clinically related, that affect patient satisfaction ratings. More research is warranted to investigate the role of hospital consumer surveys' ratings in the spine surgical patient population.

Keywords: satisfaction; spine surgery; neurosurgery; Press Ganey; HCAHPS; survey

Introduction

In recent years, patient satisfaction measures have played an increasingly important role in the evaluation of hospital and surgeon performance, quality of care, and reimbursement.^{1, 2} Although objective and functional outcomes have been widely studied in both neurosurgical and orthopedic spine operations,³ there exists a relative void in systematic assessments of this population's subjective patient satisfaction measures. Patient satisfaction can be broadly categorized as contentment with the diagnosis process, delivery of care, and/or clinical outcomes.⁴ A satisfactory clinical outcome, such as cessation of pain, may be achieved despite an unsatisfactory process of care (e.g., difficult insurance process, long clinic wait-time) or delivery of care (e.g., high cost, delay of treatment, unexpected post-operative setting or length of stay). Since many factors may influence a surgical patient's evaluation of care quality and "satisfaction", attempts to quantify this multifaceted and complex outcome measure via Likert and/or visual analog scales may be inadequate and arduous.

Nonetheless, patient satisfaction measures such as Press Ganey⁵ and Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS)⁶ surveys are widely used to query patients' perceptions of their hospital experience.⁷ The Centers for Medicare and Medicaid Services even requires all U.S. hospitals to administer HCAHPS surveys to all inpatient individuals, the results of which may directly influence physician and hospital compensation and reimbursement.⁷⁻¹⁰ Regardless of the widespread utilization of these surveys, there exists no systematic review evaluating predictors of spine surgery patient satisfaction based on the two most widely utilized surveys—Press Ganey and HCAHPS. As such, this systematic review aims to analyze the available literature of spine surgery Press Ganey and HCAHPS surveys especially highlighting predictive factors of patient satisfaction.

Materials and Methods

Database Search

This study was exempt from Institutional Review Board evaluation due to its investigation of publicly accessible literature and non-involvement of human subjects. A thorough literature search of the published English-language literature was performed in the National Library of Medicine PubMed/MEDLINE, Google Scholar, and Cochrane databases using the keywords (“spine”) AND (“press ganey” OR “HCAHPS”) from database inception to July 2020 (**Figure 1**). Each abstract was reviewed independently for relevance by three authors (BML, KG, and NJB) and another author (RS) acted as the final mediator if disagreements for inclusion occurred. Each article’s reference list was screened to identify additional articles that met our inclusion criteria for this study. Full manuscripts of the studies that met inclusion criteria were evaluated independently by two authors (BML and RS). This study was conducted in agreement with the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.¹¹

Inclusion and Exclusion Criteria

Inclusion criteria consisted of 1) English language, 2) original research, and 3) utilizing Press Ganey or HCAHPS survey in spine surgery patients. Case reports, editorials/commentaries, reviews, and preliminary data in the form of abstracts were excluded. Furthermore, studies that did not indicate a particular satisfaction survey utilized or those that utilized a different measure other than Press Ganey or HCAHPS surveys were excluded.

Data collection

The primary outcome variables were positive, negative, and neutral predictors of patient satisfaction. Secondary outcome variables consisted of the type of survey utilized, patient

satisfaction definition, sample size, clinical population, study type, publication year, journal published, and the 2018 journal impact factor (via Web of Science Group *Journal Citations Report*, Clarivate Analytics, 2018). Also, study quality of the analyzed manuscripts was assessed in relation to the level of evidence (I-IV).¹²

Statistical Analysis

Quantitative data was analyzed using descriptive statistics in the form of percentages. Top-box scores were defined as the sum of the highest points on a satisfaction scale, with a high top-box score reflecting a higher proportion of favorable ratings and a low top-box score reflecting a lower proportion of favorable ratings. Linear regression was used to determine trends in publications across years. All statistical analyses were performed in PASW Statistics 18.0 software (SPSS Inc., Chicago, IL). A p -value of < 0.05 was statistically significant.

Results

A total of 26 articles in PubMed, 1840 articles in Google Scholar, 25 articles in Cochrane, and 8 articles by examining the references of each included paper were identified. After these 1,899 articles were screened for inclusion (**Figure 1**), 24 studies met inclusion criteria for qualitative analysis. Summary tables of all the studies included in our analysis are presented in **Table 1 and Table 2**. Many of the studies were published in journals pertaining to spinal disorders (**Table 3**). Temporal trends in the number of publications published per year demonstrated a significant increase in the number of publications year after year ($p=0.04$). Most included studies retrospectively reviewed patient records and reported demographic and/or pre-operative clinical variables along with post-operative clinical variables influencing Press Ganey or HCAPHS survey measures. For most studies, patient satisfaction with hospital experience was

defined as scores of 9-10 on HCAHPS surveys and 8-11 on Press Ganey surveys. HCAHPS surveys were utilized in most studies (n=15; 62.5%) by using the top-box/perfect score metric to measure patient satisfaction (n=12; 80%) (**Table 4**). Press Ganey survey was evaluated by less studies (n=9; 37.5%) where satisfaction was most commonly measured using continuous scale metrics (n=4; 44.4%) (**Table 4**).

The included studies evaluated the relationship between patient satisfaction and variables that could be broadly categorized as follows (not mutually exclusive): a) patient demographics (n=17; 71%), b) pre- and intra-operative clinical variables (n=5; 21%), and c) post-operative factors depending on the temporal timeline of survey administration (n=8; 33.3%). Top positive predictors of patient satisfaction were patient and nursing/medical staff relationship (n=4; 17%), physician-patient relationship (n=4; 17%), managerial oversight of received care (n=3; 13%), same sex/ethnicity between patient and physician (n=2; 8%), and older age (n=2; 8%) (**Table 5**). Top negative predictors of patient satisfaction were high Charlson Comorbidity index/high disability/worse overall health functioning (n=7; 29%), increased length of hospital stay (n=4; 17%), high rating for pain/complications/readmissions (n=4; 17%), and psychosocial factors (n=3; 13%) (**Table 5**).

Discussion

To our knowledge, this is the first study to systematically review the available literature pertaining to spine surgery patient satisfaction as evaluated through the two most administered patient satisfaction surveys—Press Ganey and HCAHPS. Though two previous reviews have broadly reported on this topic,^{13, 14} this manuscript presents a comprehensive review and analysis of the two most widely utilized patient satisfaction surveys in order to identify factors most

predictive of patient satisfaction following spine surgery. These relatively subjective measures used to quantify patient satisfaction are increasingly being utilized in hospital systems year over year.¹⁵ Therefore, it is imperative to examine predictors of positive and negative patient satisfaction to adjust practice patterns where needed. Time and continuous research will tell whether these measures can be valuable in changing delivery of care or eventual patient clinical outcomes.

Press Ganey

The Press Ganey (Nashville, Tennessee) survey was developed in 1985 by Irwin Press, PhD and Rod Ganey, PhD, and is now nationally recognized as a premier patient satisfaction survey. Due to their meticulously designed patient-centered and broad care delivery questions, there are currently greater than 20,000 hospitals and health centers utilizing this platform to evaluate patient satisfaction.¹⁶⁻¹⁸ The main criticisms with the Press Ganey survey are its large heterogeneity and ambiguity in terms of assessing and quantifying satisfaction.⁷ Olivero and colleagues recently brought into question the usefulness of Press Ganey in spine surgery patients by evaluating the association between Press Ganey scores and lumbar spine quality outcome measures from the National Neurosurgery Quality and Outcomes Database (N2QOD).¹⁹ The authors demonstrated a negative correlation between top-box Press Ganey scores and the N2QOD results. Specifically, surgeons with lower top-box Press Ganey scores had lower post-operative hospital stay lengths and complication rates.¹⁹ This negative correlation observed in their study (correlation coefficient = -0.9), therefore, demonstrates that patient satisfaction measures cannot be used as a global variable to assess the relationship between clinical outcomes and high patient satisfaction. Moreover, a recent study demonstrated that patient

satisfaction with physician communication is not necessarily correlated with post-operative patient reported health-status measures and outcomes 1-year following spine surgery.²⁰

Many studies in our review identified several key factors predictive of negative patient satisfaction that were consistent across multiple investigations, with spine patients typically reporting lower patient satisfaction than non-spine patients.²¹ This trend may suggest that achieving the goal of high patient satisfaction may not be entirely reflective of achieving “optimal overall care/clinical outcomes”. Instead, high patient satisfaction may be measuring an absence of patient-perceived dissatisfaction related to many potential factors (i.e., clinically and non-clinically related) as part of the care process, which is more than likely outside the control of any single provider. From our systematic review, increased post-surgical length of stay was among many factors most likely predictive of low patient satisfaction.^{19, 22} However, shorter post-surgical lengths of stay did not necessarily correlate to high top-box Press Ganey scores.²²

It is possible that having pre-existing medical conditions can impact patients’ top-box Press Ganey scores. A recent study found that Charlson/Deyo Comorbidity Index was negatively associated with receiving a high top-box Press Ganey score.²² Additionally, a study by Mazur and colleagues identified that patients with self-reported “severe” disability tended to register low top-box Press Ganey scores.²³ The potential association between comorbidities/disabilities and patient satisfaction has also been discussed in other spine surgery studies using a standard survey²⁴ or the Musculoskeletal Outcomes Data Evaluation and Management System's satisfaction survey.²⁵ Lastly, mental health such as psychological distress has been shown to negatively affect patient satisfaction. A study by Abtahi and colleagues demonstrated that “distressed-depressive” and “distressed-somatic” patients reported significantly lower overall and provider-specific satisfaction scores compared to “normal” patients, as defined by the

study's assessments.²⁶ Taken together, these studies suggest that the physical and overall mental health of patients undergoing spine surgery may influence their reported Press Ganey satisfaction scores, although one study did not find an association between overall health functioning and patient satisfaction.²⁷ As such, this can be considered when clinicians and staff are providing various services and/or communicating post-operative care to patients that are less mentally or physically capable. Also, other variables that were notably found to be associated with lower Press Ganey patient satisfaction scores were higher education level, paper/pen survey administration, having commercial/private insurance, and patients advised against a surgical operation.^{23, 28}

Moreover, there may exist several clinical and/or demographic factors associated with receiving high patient satisfaction scores that are uncontrollable by the physicians' clinical expertise and/or provided care, such as patients' older age and patients being the same sex/race as the provider.^{28, 29} On the other hand, we found that factors relating to the physician-patient relationship can potentially be associated with patient satisfaction, such as quality of physician counseling and prompt meetings based on their scheduled appointments.³⁰ Overall, these studies suggest that patient satisfaction is multifaceted and many aspects either directly controllable or uncontrollable by the clinician and/or medical staff can impact patients' perceived satisfaction or dissatisfaction.

HCAHPS

HCAHPS, produced by the Centers for Medicare & Medicaid Services and the Agency for Healthcare Research and Quality, was approved in 2005 for methodical appraisals of patient satisfaction. The survey is highly connected to hospital reimbursements through the Inpatient

Prospective Payment System and the Value-Based Purchasing program.³¹ The HCAHPS survey, which can be distributed through multiple venues (e.g., mail, telephone) and varying languages, quantifies satisfaction matching the following criteria: 1) “communication with doctors”, 2) “communication with nurses”. 3) “responsiveness of hospital staff”, 4) “pain management”, 5) “communication about medicines”, 6) “discharge information”, 7) “care transition”, 8) “cleanliness of hospital environment”, 9) “quietness of hospital environment”, 10) “recommend the hospital”, and 11) “overall hospital rating”³². Survey responses depend on the specific question and comprise of either: 1) “never, sometimes, usually, and always”, 2) “definitely no, probably no, probably yes, and definitely yes”, and 3) “strongly disagree, disagree, agree, or strongly agree”, and 4) ordinal scale of 1-10 for hospital rating. Top-box rating is most designated as the highest remark within each category: “always”, “definitely yes”, “strongly agree”, or top-box score of 9-10.

From our systematic review of the spine surgery literature, several factors predictive of both patient satisfaction and dissatisfaction measured through the HCAHPS survey were identified. The association between HCAHPS-measured satisfaction and patient demographic and preoperative factors were investigated in three studies pertaining to spine surgical patients.³³⁻
³⁶ It was observed that allergies or depression in patients negatively impacted their reported satisfaction.^{33, 34} Specifically, Levin and colleagues observed that pre-operative allergies were negatively associated with communication between patients and the nursing staff, pain management, communication about post-operative medicine use and medical care.³⁴ Two separate studies by Levin and colleagues observed that preoperative depression was negatively associated with satisfaction pertaining to doctor/nursing communication with patients, hospital staff alertness, discharge communication, overall rating of a hospital,³³ and doctor

respect/listening³⁷ in the lumbar and cervical spine surgical patient populations, respectively. Additionally, following a prospective study of 200 spine clinic patients, Bible and colleagues found that less formal education, younger age, and smoking were associated with lower patient satisfaction.³⁵ These studies further illustrate how factors objectively distinct from the physician's surgical expertise and clinical outcomes may affect the patient's perceived satisfaction following spine surgical operations.

Moreover, other spine surgical studies have attempted to uncover independent predictors for patient satisfaction assessed through HCAHPS. In a retrospective study on 1480 spine surgery patients, Mets and colleagues observed that top-box ratings were associated with older age, male gender, and undergoing cervical (compared to lumbar) spine surgery, while non top-box ratings were associated with worse overall health, undergoing non-elective procedures, increased post-operative length of stay, and experiencing any complication or requiring hospital readmission.³⁸ In another study of 391 lumbar spine surgery patients, longer post-operative length of stay was associated with lower top-box satisfaction for doctor listening and pain management subdivisions of the HCAHPS survey.³⁹ The role of post-operative readmissions was also assessed in a separate study of 453 lumbar spine surgery patients, where increased volume of post-discharge emergency department visits was associated with lower top-box overall scores for doctor respect/listening and discharge information subdivisions of the HCAHPS survey.⁴⁰ Additionally, Hopkins and colleagues' study of 1118 spine patients also noted lower top-box HCAHPS scores for patients with increased post-operative length of stay, lower overall health status, and patients undergoing revision surgeries, while no positive predictors of patient satisfaction were observed, suggesting that patient satisfaction may not be accurately assessed via the current measures.⁴¹ This is not an unassuming conclusion since more than half (53%) of

our HCAHPS studies did not identify positive predictors of patient satisfaction.^{33, 34, 37, 39-43} Lastly, it has been previously reported that in cases of low response rates, HCAHPS responders may not be representative of the total patient sample studied contributing to so-called nonresponder bias.⁴⁴

In addition to studies by Levin et al. and Smith et al., patient perceived satisfaction regarding pain management was also assessed in Maher and colleagues study of 562 spine surgery patients, demonstrating an association between increased total intraoperative opioid dosage and a lower top-box overall satisfaction and hospital recommendation to family and friends.⁴³ They observed a lower top-box satisfaction for pain management for patients with increased surgery length, post-operative length of stay, and total intraoperative Ketorolac dosage. Additionally, Kerezoudis and colleagues observed that spine surgical patients reported lower top-box scores for pain management.⁴⁵ Hence, in addition to the surgical operation length and post-operative length of stay, patients' expectations or understanding thereof regarding pain management is important to consider as it may affect perceived patient satisfaction.

Moreover, some of the HCAHPS studies shed light on the potential influence of hospital staff engagement with patients during and following the surgical operation. For example, Levin and colleagues' study of 453 lumbar spine surgery patients observed that receipt of a top-box overall score was more often observed for patients with high satisfaction in the following HCAHPS subdivisions: a) doctor medication description, b) doctors and/or nursing staff respectfulness, c) doctor and/or nursing staff perceived listening proficiencies, d) description of medication side effects, e) regard for personal and family inclinations, f) facility cleanliness/hygiene and quietness, g) patient comprehension of post-operative care, and h) pain management.⁴⁶ A separate study by Bible and colleagues showed that perceived provider

compassion, along with swift appointment arranging, staff teamwork, nurse/resident engagement, wait times, and post-operative medical communications were highly associated with positive satisfaction results.⁴⁷ In attempts to understand factors that could improve the nursing/medical staff scores, Wang et al.'s study of 741 spine surgery patients reported that nurse/medical staff explanation scores were enhanced for patients who received goal setting and smart-room technology educational videos in comparison to those solely receiving their discharge time/date via written communication.⁴⁸ These studies suggest that aspects pertinent to interactions and communications between patients and their providers may largely affect global hospital ratings in the spine surgery patient population as measured through HCAHPS.

Limitations

Although we took great care in the collection and analysis of this data, there are several distinct limitations with our current study. First, our current conclusions regarding potential predictors of patient satisfaction is limited to our low sample size, and there exists large heterogeneity in our analyses of the included studies. Second, given the nature of the data reporting, we were unable to make reasonable comparisons between the HCAHPS and Press Ganey survey measures. Therefore, our analysis was segregated into separate descriptive and pooled analyses. Regardless, this manuscript reviewing all studies that evaluated patient satisfaction in spine surgery using HCAHPS and Press Ganey surveys adds an additional layer to the already multifaceted nature of understanding patient-perceived satisfaction in the spine surgery population. Continuous research in this field will help elucidate whether implementing HCAHPS and/or Press Ganey measures affect delivery of care and clinical outcomes.

Conclusion

There is much variability in the literature in terms of different factors that affect patient satisfaction. We found that positive predictors of patient satisfaction tended to be related to the relationships between patients and their providers/associated medical staff members, while negative predictors of patient satisfaction tended to be either demographic factors not controllable by the provider and/or post-operative care management. More research is warranted to investigate the role of hospital consumer surveys' ratings in the spine surgical patient population.

References

1. Farley H, Enguidanos ER, Coletti CM, et al. Patient satisfaction surveys and quality of care: an information paper. *Ann Emerg Med*. 2014;64(4): 351-357. <https://doi.org/10.1016/j.annemergmed.2014.02.021>.
2. Edwards ST, Bitton A, Hong J, Landon BE. Patient-centered medical home initiatives expanded in 2009-13: providers, patients, and payment incentives increased. *Health Aff (Millwood)*. 2014;33(10): 1823-1831. <https://doi.org/10.1377/hlthaff.2014.0351>.
3. Haws BE, Khechen B, Bawa MS, et al. The Patient-Reported Outcomes Measurement Information System in spine surgery: a systematic review. *J Neurosurg Spine*. 2019;30(3): 405-413. <https://doi.org/10.3171/2018.8.SPINE18608>.
4. Donabedian A. The quality of care. How can it be assessed? *JAMA*. 1988;260(12): 1743-1748. <https://doi.org/10.1001/jama.260.12.1743>.
5. Presson AP, Zhang C, Abtahi AM, Kean J, Hung M, Tyser AR. Psychometric properties of the Press Ganey(R) Outpatient Medical Practice Survey. *Health Qual Life Outcomes*. 2017;15(1): 32. <https://doi.org/10.1186/s12955-017-0610-3>.
6. Giordano LA, Elliott MN, Goldstein E, Lehrman WG, Spencer PA. Development, implementation, and public reporting of the HCAHPS survey. *Med Care Res Rev*. 2010;67(1): 27-37. <https://doi.org/10.1177/1077558709341065>.
7. Zusman EE. HCAHPS replaces Press Ganey survey as quality measure for patient hospital experience. *Neurosurgery*. 2012;71(2): N21-24. <https://doi.org/10.1227/01.neu.0000417536.07871.ed>.
8. HCAHPS: Patients' Perspectives of Care Survey; 2019. <https://www.cms.gov/Medicare/Quality-Initiatives-Patient-Assessment-Instruments/HospitalQualityInits/HospitalHCAHPS.html>.
9. Elliott MN, Beckett MK, Lehrman WG, et al. Understanding The Role Played By Medicare's Patient Experience Points System In Hospital Reimbursement. *Health Aff (Millwood)*. 2016;35(9): 1673-1680. <https://doi.org/10.1377/hlthaff.2015.0691>.
10. Zamora D. Using patient satisfaction as a basis for reimbursement: political, financial, and philosophical implications. *Creat Nurs*. 2012;18(3): 118-123.
11. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med*. 2009;6(7): e1000097. <https://doi.org/10.1371/journal.pmed.1000097>.
12. Yarascavitch BA, Chuback JE, Almenawer SA, Reddy K, Bhandari M. Levels of evidence in the neurosurgical literature: more tribulations than trials. *Neurosurgery*. 2012;71(6): 1131-1137; discussion 1137-1138. <https://doi.org/10.1227/NEU.0b013e318271bc99>.
13. Malpani R, Hilibrand, A.S., Grauer, J.N. Evolution and Use of Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) Surveys and Their Application for Spinal Surgery Patients. *Contemporary Spine Surgery*. 2018;19(5): 8. <https://doi.org/10.1097/01.CSS.0000532433.06327.f4>.
14. Menendez JY, Omar NB, Chagoya G, et al. Patient Satisfaction in Spine Surgery: A Systematic Review of the Literature. *Asian Spine J*. 2019. <https://doi.org/10.31616/asj.2019.0032>.

15. Elliott MN, Cohea CW, Lehrman WG, et al. Accelerating Improvement and Narrowing Gaps: Trends in Patients' Experiences with Hospital Care Reflected in HCAHPS Public Reporting. *Health Serv Res.* 2015;50(6): 1850-1867. <https://doi.org/10.1111/1475-6773.12305>.
16. Urden LD. Patient satisfaction measurement: current issues and implications. *Outcomes Manag.* 2002;6(3): 125-131.
17. Rogers F, Horst M, To T, et al. Factors associated with patient satisfaction scores for physician care in trauma patients. *J Trauma Acute Care Surg.* 2013;75(1): 110-114; discussion 114-115. <https://doi.org/10.1097/TA.0b013e318298484f>.
18. Boulding W, Glickman SW, Manary MP, Schulman KA, Staelin R. Relationship between patient satisfaction with inpatient care and hospital readmission within 30 days. *Am J Manag Care.* 2011;17(1): 41-48.
19. Olivero W, Wang H, Vinson D, Pierce C, Trumbull S. Correlation Between Press Ganey Scores and Quality Outcomes From The National Neurosurgery Quality and Outcomes Database (Lumbar Spine) for a Hospital Employed Neurosurgical Practice. *Neurosurgery.* 2018;65(CN_suppl_1): 34-36. <https://doi.org/10.1093/neuros/nyy106>.
20. Rabah NM, Levin JM, Winkelman RD, Mroz TE, Steinmetz MP. The Association Between Physicians' Communication and Patient Reported Outcomes in Spine Surgery. *Spine (Phila Pa 1976).* 2020. <https://doi.org/10.1097/BRS.0000000000003458>.
21. Chen YR, Johnson E, Montalvo C, et al. Patient Satisfaction and Press Ganey Scores for Spine Versus Nonspine Neurosurgery Clinics. *Clin Spine Surg.* 2019;32(4): E188-E192. <https://doi.org/10.1097/BSD.0000000000000825>.
22. Hopkins BS, Patel MR, Yamaguchi JT, Cloney MB, Dahdaleh NS. Predictors of patient satisfaction and survey participation after spine surgery: a retrospective review of 17,853 consecutive spinal patients from a single academic institution. Part 1: Press Ganey. *J Neurosurg Spine.* 2019;30(3): 382-388. <https://doi.org/10.3171/2018.8.SPINE18594>.
23. Mazur MD, McEvoy S, Schmidt MH, Bisson EF. High self-assessment of disability and the surgeon's recommendation against surgical intervention may negatively impact satisfaction scores in patients with spinal disorders. *J Neurosurg Spine.* 2015;22(6): 666-671. <https://doi.org/10.3171/2014.10.SPINE14264>.
24. Yamashita K, Hayashi J, Ohzono K, Hiroshima K. Correlation of patient satisfaction with symptom severity and walking ability after surgical treatment for degenerative lumbar spinal stenosis. *Spine (Phila Pa 1976).* 2003;28(21): 2477-2481. <https://doi.org/10.1097/01.BRS.0000090888.63860.4F>.
25. Soroceanu A, Ching A, Abdu W, McGuire K. Relationship between preoperative expectations, satisfaction, and functional outcomes in patients undergoing lumbar and cervical spine surgery: a multicenter study. *Spine (Phila Pa 1976).* 2012;37(2): E103-108. <https://doi.org/10.1097/BRS.0b013e3182245c1f>.
26. Abtahi AM, Brodke DS, Lawrence BD, Zhang C, Spiker WR. Association between patient-reported measures of psychological distress and patient satisfaction scores after spine surgery. *J Bone Joint Surg Am.* 2015;97(10): 824-828. <https://doi.org/10.2106/JBJS.N.00916>.
27. Abtahi AM, Lyman KS, Brodke DS, Lawrence BD, Zhang C, Spiker WR. Patient Satisfaction is Not Associated With Self-reported Disability in a Spine Patient Population. *Clin Spine Surg.* 2017;30(8): E1165-E1168. <https://doi.org/10.1097/BSD.0000000000000431>.

28. Johnson BC, Vasquez-Montes D, Steinmetz L, et al. Association Between Nonmodifiable Demographic Factors and Patient Satisfaction Scores in Spine Surgery Clinics. *Orthopedics*. 2019;42(3): 143-148. <https://doi.org/10.3928/01477447-20190424-05>.
29. Steinmetz L, Vasquez-Montes D, Johnson BC, et al. Modifiable and nonmodifiable factors associated with patient satisfaction in spine surgery and other orthopaedic subspecialties: A retrospective survey analysis. *Current Orthopaedic Practice*. 2019;30(6): 555-560. <https://doi.org/10.1097/bco.0000000000000800>.
30. Etier BE, Jr., Orr SP, Antonetti J, Thomas SB, Theiss SM. Factors impacting Press Ganey patient satisfaction scores in orthopedic surgery spine clinic. *Spine J*. 2016;16(11): 1285-1289. <https://doi.org/10.1016/j.spinee.2016.04.007>.
31. Tevis SE, Kennedy GD, Kent KC. Is There a Relationship Between Patient Satisfaction and Favorable Surgical Outcomes? *Adv Surg*. 2015;49: 221-233. <https://doi.org/10.1016/j.yasu.2015.03.006>.
32. Services DoHaHSCfMM. CAHPS® Hospital Survey (HCAHPS) Quality Assurance Guidelines Version 12.0. Baltimore, MD: Centers for Medicare & Medicaid Services; 2017.
33. Levin JM, Winkelman RD, Smith GA, et al. Impact of Preoperative Depression on Hospital Consumer Assessment of Healthcare Providers and Systems Survey Results in a Lumbar Fusion Population. *Spine (Phila Pa 1976)*. 2017;42(9): 675-681. <https://doi.org/10.1097/BRS.0000000000002101>.
34. Levin JM, Boyle S, Winkelman RD, et al. Patient-reported Allergies are Associated With Preoperative Psychological Distress and Less Satisfying Patient Experience in a Lumbar Spine Surgery Population. *Clin Spine Surg*. 2018;31(7): E368-E374. <https://doi.org/10.1097/BSD.0000000000000665>.
35. Bible JE, Kay HF, Shau DN, O'Neill KR, Segebarth PB, Devin CJ. What Patient Characteristics Could Potentially Affect Patient Satisfaction Scores During Spine Clinic? *Spine (Phila Pa 1976)*. 2015;40(13): 1039-1044. <https://doi.org/10.1097/BRS.0000000000000912>.
36. Galivanche AR, Mercier MR, Adrados M, et al. Admission NarxCare Narcotics Scores are not Associated with Adverse Surgical Outcomes or Self-Reported Patient Satisfaction Following Elective Spine Surgery. *Spine (Phila Pa 1976)*. 2019. <https://doi.org/10.1097/BRS.00000000000003120>.
37. Levin JM, Rabah NM, Winkelman RD, Mroz TE, Steinmetz MP. The Impact of Preoperative Depression on Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) Survey Results in a Cervical Spine Surgery Setting. *Spine (Phila Pa 1976)*. 2019. <https://doi.org/10.1097/BRS.00000000000003222>.
38. Mets EJ, Mercier MR, Hilibrand AS, Scott MC, Varthi AG, Grauer JN. Patient-Related Factors and Perioperative Outcomes Are Associated with Self-Reported Hospital Rating after Spine Surgery. *Clin Orthop Relat Res*. 2019. <https://doi.org/10.1097/CORR.0000000000000892>.
39. Smith GA, Chirieleison S, Levin J, et al. Impact of length of stay on HCAHPS scores following lumbar spine surgery. *J Neurosurg Spine*. 2019: 1-6. <https://doi.org/10.3171/2019.3.SPINE181180>.
40. Levin JM, Winkelman RD, Smith GA, et al. Emergency department visits after lumbar spine surgery are associated with lower Hospital Consumer Assessment of Healthcare Providers and Systems scores. *Spine J*. 2018;18(2): 226-233. <https://doi.org/10.1016/j.spinee.2017.06.043>.

41. Hopkins BS, Patel MR, Yamaguchi JT, Cloney MB, Dahdaleh NS. Predictors of patient satisfaction and survey participation after spine surgery: a retrospective review of 17,853 consecutive spinal patients from a single academic institution. Part 2: HCAHPS. *J Neurosurg Spine*. 2019;30(3): 389-396. <https://doi.org/10.3171/2018.8.SPINE181024>.
42. Levin JM, Winkelman RD, Smith GA, et al. The association between the Hospital Consumer Assessment of Healthcare Providers and Systems (HCAHPS) survey and real-world clinical outcomes in lumbar spine surgery. *Spine J*. 2017;17(11): 1586-1593. <https://doi.org/10.1016/j.spinee.2017.05.002>.
43. Maher DP, Wong W, Woo P, et al. Perioperative factors associated with HCAHPS responses of 2,758 surgical patients. *Pain Med*. 2015;16(4): 791-801. <https://doi.org/10.1111/pme.12651>.
44. Malpani R, Adrados M, Mercier MR, et al. Characteristics and Predictors of HCAHPS Nonresponse After Spine Surgery. *Spine (Phila Pa 1976)*. 2020;45(8): E448-E456. <https://doi.org/10.1097/BRS.0000000000003287>.
45. Kerezoudis P, Alvi MA, Ubl DS, et al. The impact of spine disease, relative to cranial disease, on perception of health and care experience: an analysis of 1484 patients in a tertiary center. *J Neurosurg*. 2018;129(6): 1630-1640. <https://doi.org/10.3171/2017.7.JNS17991>.
46. Levin JM, Winkelman RD, Tanenbaum JE, Benzel EC, Mroz TE, Steinmetz MP. Key drivers of patient satisfaction in lumbar spine surgery. *J Neurosurg Spine*. 2018;28(6): 586-592. <https://doi.org/10.3171/2017.10.SPINE17732>.
47. Bible JE, Shau DN, Kay HF, Cheng JS, Aaronson OS, Devin CJ. Are Low Patient Satisfaction Scores Always Due to the Provider?: Determinants of Patient Satisfaction Scores During Spine Clinic Visits. *Spine (Phila Pa 1976)*. 2018;43(1): 58-64. <https://doi.org/10.1097/BRS.0000000000001453>.
48. Wang W, Dudjak LA, Larue EM, Ren D, Scholle C, Wolf GA. The influence of goal setting and SmartRoom patient education videos on readmission rate, length of stay, and patient satisfaction in the orthopedic spine population. *Comput Inform Nurs*. 2013;31(9): 450-456. <https://doi.org/10.1097/NXN.0b013e3182999df8>.

Figure Legends

Figure 1. PRISMA study selection flow diagram.

Table 1: Included Study Characteristics						
Citation	Journal Name	Study Type (Retrospective vs Prospective)	Sample Size	Clinical Patient Population	Assessment (Press Ganey vs HCAPHS)	Definition of Patient Satisfaction
Steinmetz et al., 2019	Current Orthopaedic Practice	Retrospective	1,400	Ambulatory Spine Surgery Patients	Press Ganey	Overall Provider Rating (0-10) and Likelihood to Recommend
Levin et al., 2019	Spine	Retrospective	145	Cervical Spine Surgery	HCAHPS	Top box (9-10)
Hopkins et al., 2019	Journal of Neurosurgery: Spine	Retrospective	1936	Neurosurgical or Orthopedic Spine Patients	Press Ganey	Perfect score (5/5)
Hopkins et al., 2019	Journal of Neurosurgery: Spine	Retrospective	1118	Neurosurgical or Orthopedic Spine Patients	HCAPHS	Perfect score (4/4)
Smith et al., 2019	Journal of Neurosurgery: Spine	Retrospective	391	Lumbar Spine Surgery Patients	HCAPHS	Top box (9-10)
Mets et al., 2019	Clinical Orthopaedics and Related Research	Retrospective	1480	Neurosurgical or Orthopedic Spine Patients	HCAPHS	Top box (9-10)
Galivanch e et al., 2019	Spine	Retrospective	69	Elective Spine Surgery Patients	HCAPHS	Top box (9-10)
Johnson et al., 2019	Orthopedics	Retrospective	1400	Ambulatory Spine Surgery Patients	Press Ganey	Not specified
Chen et al., 2019	Clinical Spine Surgery	Retrospective	578	Neurosurgical Spine Patients	Press Ganey	Continuous (0-100)
Olivero et al., 2018	Neurosurgery	Retrospective	77	Lumbar Spine Surgery Patients	Press Ganey	Top box (5 or 9-10)
Levin et	Clinical Spine	Retrospective	421	Lumbar Spine	HCAPHS	Top box

al., 2018	Surgery			Surgery Patients		(9-10)
Levin et al., 2018	Journal of Neurosurgery: Spine	Retrospective	453	Lumbar Spine Surgery Patients	HCAPHS	Top box (9-10)
Levin et al., 2018	The Spine Journal	Retrospective	453	Lumbar Spine Surgery Patients	HCAPHS	Top box (9-10)
Kerezoudis et al., 2018	Journal of Neurosurgery	Retrospective	1484	Elective Spine Surgery Patients	HCAPHS	Top box (9-10)
Bible et al., 2018	Spine	Prospective	158	Unspecified Spine Surgery Patients	Modified HCAPHS	Ordinal (1-10)
Levin et al., 2017	The Spine Journal	Retrospective	249	Lumbar Spine Surgery Patients	HCAPHS	Top box (9-10)
Levin et al., 2017	Spine	Retrospective	237	Lumbar Spine Fusion Surgery Patients	HCAPHS	Top box (9-10)
Abtahi et al., 2017	Clinical Spine Surgery	Retrospective	231	Unspecified Spine Surgery Patients	Press Ganey	Continuous (0-100)
Etier et al., 2016	The Spine Journal	Retrospective	353	Orthopedic Spine Surgery Patients	Press Ganey	Score of 8-11 (satisfied)
Mazur et al., 2015	Journal of Neurosurgery: Spine	Retrospective	130	Neurosurgical Spine Disorders Patient Population	Press Ganey	Continuous (0-100)
Bible et al., 2015	Spine	Prospective	200	Unspecified Spinal Disorders Patient Population	Modified HCAPHS	Ordinal (1-10)
Maher et	Pain Medicine	Retrospective	562	Unspecified	HCAPHS	Top box

al., 2015				Spine Surgery Patients		(9-10)
Abtahi et al., 2015	The Journal of Bone and Joint Surgery	Retrospective	103	Unspecified Spine Surgery Patients	Press Ganey	Continuous (0-100)
Wang et al., 2013	Computers, Informatics, Nursing	Retrospective	741	Orthopedic Spine Surgery Patients	HCAPHS	Continuous (0-100)

Table 2: Predictors of Patient Satisfaction

Citation	Predictors of High Patient Satisfaction or Negative Predictors of Low Patient Satisfaction	Predictors of Low Patient Satisfaction or Negative Predictors of High Patient Satisfaction	Comments
Steinmetz et al., 2019	Provider Communication, nursing/medical staff communication, and same ethnicity	Not identified	N/A
Levin et al., 2019	None identified	Depressed patients reported lower satisfaction for doctor respect/listening (overall doctor communication)	Depressed patients were younger in age, more likely to have preoperative neck pain, and lower quality of life
Hopkins et al., 2019	None identified	High Charlson Comorbidity Index, increased time since surgical operation/discharge, and increased post-operative hospital stay	The perceived expertise of the surgeon was not a prognosticator for top-box or low Press Ganey scores; Patients who had a posterior-approach surgical procedure in comparison to other procedures were less likely to note a low (≤ 2) Press Ganey score
Hopkins et al., 2019	None Identified	Revision surgeries, urgent/emergency procedures, increased time since surgery/discharge, increased post-operative hospital stay, and high Charlson Comorbidity Index	N/A
Smith et al., 2019	None Identified	Patients with high post-operative hospital stay were half as likely to report their physician	N/A

		either a) listened to them, or b) treated them with courtesy and respect; longer post-operative hospital stay were associated with lower scores for c) pain management, d) explanations of pain medication administration, and e) overall hospital rating	
Mets et al., 2019	Older age (>40 years), gender identification as male, and cervical surgery (compared to lumbar surgery)	American Society of Anesthesiologists (ASA) classification ≥ 2 , non-elective surgery, having any adverse event within 30 days, readmission within 30 days, and longer post-operative hospital stay	Patients' hospital ratings did not differ by surgeon's specialty (neurosurgery vs orthopedics).
Galivanche et al., 2019	Not investigated	Not investigated	HCAPHS responses were not significantly correlated with admission narcotics scores.
Johnson et al., 2019	Older age and same sex/ethnicity as provider	Higher education level (graduate level education), paper and pen/pencil survey administration, and commercial/private insurance	N/A
Chen et al., 2019	Not investigated	Not investigated	In comparison to non-spine patients, spine patients noted overall lower satisfaction, physician/nurse care, personal concerns,

			admission/visitor environments, and overall hospital rating.
Olivero et al., 2018	Low National Neurosurgery Quality and Outcomes Database (N2QOD) scores were associated with high Press Ganey scores	High National Neurosurgery Quality and Outcomes Database (N2QOD) scores were associated with lower Press Ganey scores; high volume of post-operative complications and longer post-operative hospital stay were associated with low Press Ganey scores	N/A
Levin et al., 2018	None identified	Higher quantity of patient-reported medical and environmental allergies	N/A
Levin et al., 2018	Staff managing pain, respect/listening from nursing staff/physician, communication of medication side effects, cleanliness, post-operative health care was explained by nurses/physicians, quietness, attentiveness of staff, and perceived quality of care delivered	Lower overall health status	Previous lumbar spine surgery, history of chronic renal failure, Caucasian race, history of cancer, length of post-operative hospital stay, and mental health status were not associated with top-box HCAPHS scores
Levin et al., 2018	None identified	Post-discharge emergency department visits	Prior lumbar surgery, history of coronary artery disease, in-hospital complications, length of post-operative hospital stay, and pre-operative EuroQol 5-Dimensions index score were not

			associated with top-box HCAPHS scores
Kerezoudis et al., 2018	In comparison to degenerative spine disease patients, oncology, and other disease (e.g., intracranial aneurysm, normal pressure hydrocephalus, and movement disorders) patients recorded higher top-box scores; in comparison to cranial patients, spinal patients reported higher top-box scores in delivery of discharge information	In comparison to cranial patients, spine patients recorded lower top-box scores in pain management and communication/responsiveness of doctors/ staff	N/A
Bible et al., 2018	Univariate: appointment scheduling, office staff friendliness, teamwork (nurse-practitioner/resident involvement), low wait times, provider interactions/behavior, treatment, and follow-up communication; multivariate: medical care communication, empathy, teamwork of staff, and follow-up communication	None identified	Pain medication prescribing and provider time spent with patients was not correlated with satisfaction; analysis was performed on both operative and non-operative spine cases
Levin et al., 2017	None identified	None identified	Pre-operative to 1-year post-operative increases in optimal clinical outcome measures defined as EuroQol 5 Dimensions (EQ-5D), Pain Disability

			Questionnaire (PDQ), and Visual Analog Score for back pain (VAS-BP) were not significantly correlated with improved top-box scores
Levin et al., 2017	None identified	Depressed patients were associated with recording lower top-box scores than non-depressed patients on multiple survey components: doctor/nursing communication, hospital recommendation, help received; tobacco use was associated with lower scores for physician respect	N/A
Abtahi et al., 2017	None identified	None identified	No significant correlations between the Oswestry Disability Index (ODI), the Neck Disability Index (NDI), and the EuroQol (EQ)-5D measures and patient satisfaction in new, recurring, and post-operative spine patients
Etier et al., 2016	Lower pain score, "definitely" response to "provider spent enough time with you" question on Press Ganey survey, and patients seen close to appointment scheduled time	Not investigated	N/A
Mazur et al., 2015	Not investigated	Satisfaction was lower for	Analysis was not on post-

		patients with severe disability and had received a recommendation against surgical operation	operative spine patients, but during consult visit for outpatient back pain work-up
Bible et al., 2015	Patients with Medicare and non-work-related injuries/medical care	Younger age, male, less formal education, patients treated through worker's compensation claim, and smoking history	Marital status, working status, pain/mental health status, travel distance, quantity of prior treatments, cervical versus lumbar spine surgery, and narcotic use were not correlated with satisfaction
Maher et al., 2015	None identified	Low hospital score noted for patients with increased opioid dosage, lower amount of help received post-operatively, increased intraoperative ketorolac dosage, and higher pain scores, increased hospitalization time, and longer post-anesthesia care unit stay	N/A
Abtahi et al., 2015	Not investigated	“distressed” spine patients gave lower global and provider satisfaction than their “normal” counterparts	Distress was measured by the Distress and Risk Assessment Method (DRAM) questionnaire
Wang et al., 2013	Higher satisfaction in nurse/staff explanation of medications observed for patients with goal setting and smart room technology education videos contrasted with patients receiving	None identified	The surgical flight plan is a written information patient received regarding their date and time of post-operative hospital discharge

	“surgical flight plan”		
--	------------------------	--	--

Journal Pre-proof


Table 3: Journals Publishing Spine Satisfaction		
Journal Title	N = 24	Impact Factor*
<i>Journal of Neurosurgery: Spine</i>	5	3.0
<i>Spine</i>	5	2.9
<i>The Spine Journal</i>	3	3.2
<i>Clinical Spine Surgery</i>	3	1.7
<i>The Journal of Bone and Joint Surgery</i>	1	4.7
<i>Neurosurgery</i>	1	4.6
<i>Clinical Orthopedics and Related Research</i>	1	4.2
<i>The Journal of Neurosurgery</i>	1	4.1
<i>Pain Medicine</i>	1	2.8
<i>Orthopedics</i>	1	1.6
<i>Computers, Informatics, Nursing</i>	1	1.0
<i>Current Orthopaedic Practice</i>	1	NA
*2019 Journal Citation Reports		

Table 4: Surveys used to assess patient satisfaction.	
<i>HCAHPS (N=15; 62.5%)</i>	Number of studies (%*)
Perfect Score/Top-box	12 (80.0)
Ordinal	2 (13.3)
Continuous	1 (6.7)
<i>Press Ganey (N=9; 37.5%)</i>	Number of studies (%*)
Continuous	4 (44.4)
Perfect Score/Top-box	3 (33.3)
Unspecified/Other	2 (22.2)
*Add up to 100% within HCAHPS or Press Ganey cohorts.	

Table 5: Top Positive and Negative Predictors of Patient Satisfaction		
<i>Top Positive Predictors</i>	N	%
Nursing/medical staff-patient relationship	4	17
Managerial oversight of care	3	13
Physician-patient relationship	3	13
Same sex/ethnicity between patient and physician	2	8
Older age	2	8
<i>Top Negative Predictors</i>	N	%
High Charlson Comorbidity/Disability/Overall Health	7	29
Increased Length of Stay	4	17
Pain Management/Complications/Readmissions	4	17
Psychosocial factors	3	13



PRISMA Flow Diagram



From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.